



STANZA[®] WP

Fungicide

Product Name:	Stanza WP
Type	Fungicide
Active Constituent	462g/kg Prochloraz present as the manganese chloride complex
Resistance group	Group 3 Fungicide - (DMI Fungicide)
Formulation	wettable powder
Pack Size	1kg 5kg (packed in a reusable bucket) Keep product out of direct sunlight
Mode of action	Systemic Curative & Prevention

Stanza is a strong broad spectrum fungicide that contains 462g/kg Prochloraz present as the manganese chloride complex

How does Stanza work?

Stanza is a member of the DMI chemical class with a FRAC of Group 3- Stanza works by preventing fungi from developing cell membranes. Without the membranes, cells cannot be made and thus the fungus dies.

Stanza not only prevents the crop from infection but also kills off existing spores which have germinated.

Pack Sizes Stanza

Stanza is available in the following pack sizes:

1kg

5kg (pre pack reusable non food use bucket)



Anthracnose Disease Mango



Anthracnose Disease Lettuce



Dry Bubble Disease Msuhrroms



COLIN CAMPBELL (CHEMICALS) PTY LTD

100% Australian owned & operated. Established 1940

HORTICULTURE · FRUIT COATINGS · TURF

www.campbellchemicals.com.au (02) 9725 2544



DIRECTIONS FOR USE

Crop	Disease	State	Rate	WHP	Comments
Field grown lettuce and nursery stock prior to transplantation (closed head varieties only)	Anthrachnose (Microdochium panattonianum)	All States	50g/100L applied in 800L water / ha (i.e. 400g/ha applied high volume)	7	Apply as a foliar spray on a 7 to 14 day schedule. Use the shorter interval when continual wet, cloudy conditions favour disease development.
Mangoes	Anthrachnose blossom blight (Colletotrichum gloeosporoides var. minor)	Qld, NSW, WA only	Dilute spraying: 100g/100L plus 200g/100L of a 800g/kg formulation of mancozeb Concentrate spraying : Refer to the Application section in GENERAL INSTRUCTIONS for both Stanza and mancozeb rates.	-	Apply on a 3 to 4 week schedule throughout the flowering period. Use the shorter interval when weather conditions favour disease development. Ensure thorough coverage of blossoms. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. For concentrate spraying do not apply Stanza at more than 500g/100L (i.e. at a concentration factor greater than 5x)
Mushrooms	Dry bubble (Verticillium fungicola var. fungicola), Wet bubble (Mycogone perniciososa), Cobweb disease (Hypomyces rosellus)	All States	24g/0.3m ³ of peat OR 3g/m ² of casing surface	-	Incorporation Mix with the water used to wet the peat and incorporate with the peat prior to casing. Spray application following harvest of first flush If Stanza has not been incorporated in the peat and disease becomes a problem, spray the casing immediately after harvest of the first flush. Use adequate water to thoroughly wet the casing.
Proteas	Damping-off and die-back (Colletotrichum spp)	All States (NOT ACT or NT)	300g/100L	-	Commence spraying in early spring and apply at 10 to 14 day intervals depending on disease pressure. Use the shorter interval when weather conditions favour disease development. Spray the entire plant using a sufficient volume to ensure thorough coverage (just before point of run-off).
Violas	Anthrachnose (Colletotrichum violae-tricoloris), Spot anthracnose (Sphaceloma violae)	All States	300g/100L	-	Spray at 7 to 28 day intervals. Use the shorter interval when conditions favour disease development.