

Tridim[®] 250EC

FUNGICIDE

GROUP 3 FUNGICIDE

Active Constituent: 250g/L TRAUDIMENOL

Poison Schedule: S6 (POISON)

APVMA Approval No: 51580



Eligible containers
5L

For systemic control of ring spot on cabbage, cauliflower and broccoli, powdery mildew on cucurbits, grapevines and papaws, pineapple disease of sugar cane and various diseases of turf as indicated in the directions for use table.

Pack Size:

5L (4 x 5L per carton)
1L (12 x 1L per carton)

Dangerous Goods Class:

Not classified as a dangerous good under the Australian Code for Transport and Storage of Dangerous Goods in Australia.

DIRECTIONS FOR USE				
Restraint: Do not apply to turf earlier than 5 weeks after sowing.				
Crop	Disease	State	Rate	Critical Comments
Cabbage Cauliflower Broccoli	Ring Spot	All states	200 or 400 mL/ha plus a non-ionic wetter 200 mL/ha	Apply every 10 days. Start the program immediately following the first two consecutive days of rain after transplanting. Use higher rate under heavy disease pressure or when weather conditions favour development of the disease.
Cucurbits	Powdery mildew	All States	40 mL/100L or 400 mL/ha	Apply a non-DMI fungicide on a schedule commencing as soon as possible after emergence until fruit set. Then either tank mix or alternate Tridim with a non-DMI fungicide on a 5-10 day schedule. Use short spray intervals when disease pressure is higher plants are growing rapidly. Good coverage is essential for good results.
Grapevines	Powdery mildew	All States	10 mL/100L or 100 mL/ha	Apply thoroughly in the following 5 spray program: 1- when shoots are 10 cm long. 2- pre-flowering. 3- at flowering. 4- after fruit set. 5- before bunches close. Do not apply in volumes less than 250 L/ha. For spray volumes between 250 and 1000L/ha, use Tridim at 100 mL/ha. For volume greater than 1000L/ha, use the volumerate of 10 mL/100L.

Tridim[®] 250EC

DIRECTIONS FOR USE

Restraint: Do not apply to turf earlier than 5 weeks after sowing.

Crop	Disease	State	Rate	Critical Comments
Papaws	Powdery mildew	QLD, NSW, WA only	20 mL/100L or 200 mL/ha	Spray when powdery mildew infection begins or when conditions favour the disease. Repeat application every 14 days while conditions favour the disease. Ensure thorough coverage of foliage particularly the crown region
Sugar cane	Pineapple disease	QLD, NSW only	20 mL/100L	Apply to setts by dipping or spraying. Ensure thorough wetting of cut ends.
Turf	Dollar spot	QLD, NSW, VIC, SA, WA only	Preventive 30mL/100m ²	Apply thoroughly on a monthly programme starting immediately the disease appears.
			Curative 60mL/100m ²	Apply thoroughly under conditions of heavy disease pressure then treat as necessary.
	Black Helminthosporium	QLD, NSW, TAS only	60mL/100m ²	Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Apply during period of turf growth on a monthly programme.
	Take – all patch	QLD, NSW, VIC, TAS, WA only		Preventive only after soil fumigation with methyl bromide. Apply thoroughly, using high volume equipment, 6-8 weeks after seeding. Repeat applications (every 8 weeks in the first two years) for continued protection.
Winter Fusarium	NSW, VIC, TAS only		Apply thoroughly on a monthly programme from end of March until danger of infection has passed.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER,
CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER
APPROPRIATE LEGISLATION.

Tridim[®] 250EC

WITHHOLDING PERIODS:

CUCURBITS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION. CABBAGE, CAULIFLOWER, BROCCOLI , GRAPEVINES, PAPAWS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION. CABBAGE, CAULIFLOWER, BROCCOLI: DO NOT GRAZE ANY TREATED AREA, OR CUT FOR STOCK FOOD.

GENERAL INSTRUCTIONS

Tridim 250 EC is a broad spectrum, systemic, foliar fungicide that is quickly taken up by plants and translocated within the plant. Because of its systemic activity, Tridim is less affected by rainfall or irrigation than contact type fungicides.

MIXING

Half fill the spray tank or vat with clean water. Then add the required amount of Tridim with agitator going and top up the tank with water. Agitate thoroughly prior to and during spraying.

GROUP	3	FUNGICIDE
-------	---	-----------

FUNGICIDE RESISTANCE WARNING

Tridim is a member of the DMI group of fungicides. For fungicide resistance management Tridim is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Tridim and other Group 3 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Tridim and other Group 3 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Colin Campbell (Chemicals) Pty, Ltd., accepts no liability for any losses that may result from the failure of Tridim to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Harmful to fish. Do not contaminate streams, rivers or waterways with this product or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor of Poisons Information Centre. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. Further information refer to the Material Safety Data Sheet MSDS 51.

CONDITIONS OF SALE

Agricultural, horticultural and pastoral preparations in their application involve varying factors such as differing conditions, soil, climate and methods of application over which the vendor does not have control, Whilst Colin Campbell (Chemicals) Pty. Ltd believes that all goods sold by it are true to label and are effective and safe for purpose indicated the company and the seller hereby expressly negate and exclude any express or implied condition, statement or warranty, statutory or otherwise, as to quality or fitness of any goods sold for any purpose or purposes whatsoever except such warranties and conditions, if any, as are implied by the Trade Practices Act 1974 (Commonwealth). The company and the seller accept no responsibility for any loss, harm or damage whatsoever suffered from the use of such goods for any purpose or purposes irrespective of whether or not the buyer was acting in reliance upon the advice recommendation or representation of the seller or any representative agent of employee of the company as to such use except in respect of breaches of conditions and warranties, if any, implied by the Trade practices Act and in respect of such breaches the liability of the company and the seller shall be limited to the replacement of the goods or the supply of equivalent goods, or the payment of the cost of replacing the goods.

APVMA Approval No: 51580

®Tridim is a registered trademark of Colin Campbell (Chemicals) Pty Ltd